























9. Cheng C-C, Ko J-Y. Early reduction for congenital dislocation of the knee within twenty-four hours of birth. *Chang Gung Med J.* 2010;33(3):266-273.
10. Mehrafshan M, Wicart P, Ramanoudjame M, Seringe R, Glorion C, Rampal V. Congenital dislocation of the knee at birth – Part I: Clinical signs and classification. *Orthop Traumatol Surg Res.* 2016;102(5):631-633. doi:10.1016/j.otsr.2016.04.008
11. Bell M, Atkins R, Sharrard W. Irreducible congenital dislocation of the knee. Aetiology and management. *J Bone Jt Surg.* 1987;69-B(3):403-406. doi:10.1302/0301-620x.69b3.3584194
12. Roy DR, Crawford AH. Percutaneous quadriceps recession: a technique for management of congenital hyperextension deformities of the knee in the neonate. *J Pediatr Orthop.* 1989;9(6):717-719.
13. Shah NR, Limpaphayom N, Dobbs MB. A minimally invasive treatment protocol for the congenital dislocation of the knee. *J Pediatr Orthop.* 2009;29(7):720-725. doi:10.1097/BPO.0b013e3181b7694d
14. Patwardhan S, Shah K, Shyam A, Sancheti P. Assessment of clinical outcome of percutaneous needle quadriceps tenotomy in the treatment of congenital knee dislocation. *Int Orthop.* 2015;39(8):1587-1592. doi:10.1007/s00264-015-2806-7
15. Youssef AO. Limited Open Quadriceps Release for Treatment of Congenital Dislocation of the Knee. *J Pediatr Orthop.* 2017;37(3):192-198. doi:10.1097/BPO.0000000000000612
16. Tercier S, Shah H, Joseph B. Quadricepsplasty for congenital dislocation of the knee and congenital quadriceps contracture. *J Child Orthop.* 2012;6(5):397-410. doi:10.1007/s11832-012-0437-8
17. B K AR, Singh KA, Shah H. Surgical management of the congenital dislocation of the knee and hip in children presented after six months of age. *Int Orthop.* 2020;44(12):2635-2644. doi:10.1007/s00264-020-04759-8
18. Sud A, Chaudhry A, Mehtani A, Tiwari A, Sharma D. Functional outcome following quadriceps tendon lengthening in congenital dislocation of the knee, with special reference to extensor weakness. *Strateg Trauma Limb Reconstr.* 2009;4(3):123-127. doi:10.1007/s11751-009-0072-1
19. Oetgen ME, Walick KS, Tulchin K, Karol LA, Johnston CE. Functional results after surgical treatment for congenital knee dislocation. *J Pediatr Orthop.* 2010;30(3):216-223. doi:10.1097/BPO.0b013e3181d48375
20. Johnston CE. Simultaneous open reduction of ipsilateral congenital dislocation of the hip and knee assisted by femoral diaphyseal shortening. *J Pediatr Orthop.* 2011;31(7):732-740. doi:10.1097/BPO.0b013e31822f1b24
21. Sud A, Kumar N, Mehtani A. Femoral shortening in the congenital dislocation of the knee joint: results of mid-term follow-up. *J Pediatr Orthop Part B.* 2013;22(5):440-444. doi:10.1097/BPB.0b013e328362b8c4