

















- to effective treatment and quality of life after 2 years: A randomized controlled trial. *Spine (Phila Pa 1976)*. 2021 Jan 1;46(1):9-16.
29. Zhang Y, Liang J, Xu N, et al. 3D-printed brace in the treatment of adolescent idiopathic scoliosis: A study protocol of a prospective randomised controlled trial. *BMJ Open*. 2020;10(11):1-6.
30. Minsk MK, Venuti KD, Daumit GL, et al. Effectiveness of the Rigo Chêneau versus Boston-style orthoses for adolescent idiopathic scoliosis: a retrospective study. *Scoliosis Spinal Disord*. 2017 Mar 20;12:7.
31. Cheung PWH, Cheung JPY. Sanders stage 7b: Using the appearance of the ulnar physis improves decision-making for brace weaning in patients with adolescent idiopathic scoliosis. *Bone Joint J*. 2021 Jan;103-B(1):141-147.
32. El-Hawary R, Chukwunyerena C. Update on evaluation and treatment of scoliosis. *Pediatr Clin North Am*. 2014 Dec;61(6):1223-41.
33. Goodbody CM, Asztalos IB, Sankar WN, et al. It's not just the big kids: both high and low BMI impact bracing success for adolescent idiopathic scoliosis. *J Child Orthop*. 2016 Oct;10(5):395-404.
34. Karol LA. Effectiveness of bracing in male patients with idiopathic scoliosis. *Spine (Phila Pa 1976)*. 2001 Sep 15;26(18):2001-5.
35. Berdishevsky H, Lebel VA, Bettany-Saltikov J, et al. Physiotherapy scoliosis-specific exercises - a comprehensive review of seven major schools. *Scoliosis Spinal Disord*. 2016 Aug 4;11:20.
36. Kwan KYH, Cheng ACS, Koh HY, et al. Effectiveness of Schroth exercises during bracing in adolescent idiopathic scoliosis: results from a preliminary study-SOSORT Award 2017 Winner. *Scoliosis Spinal Disord*. 2017 Oct 16;12:32.
37. Romano M, Minozzi S, Bettany-Saltikov J, et al. Exercises for adolescent idiopathic scoliosis. *Cochrane Database Syst Rev*. 2012 Aug 15;2012(8):CD007837.
38. Williams MA, Heine PJ, Williamson EM, et al. Active Treatment for Idiopathic Adolescent Scoliosis (ACTivATeS): a feasibility study. *Health Technol Assess*. 2015 Jul;19(55):1-242.
39. Monticone M, Ambrosini E, Cazzaniga D, et al. Active self-correction and task-oriented exercises reduce spinal deformity and improve quality of life in subjects with mild adolescent idiopathic scoliosis. Results of a randomised controlled trial. *Eur Spine J*. 2014 Jun;23(6):1204-14.
40. Kuru T, Yeldan İ, Dereli EE, et al. The efficacy of three-dimensional Schroth exercises in adolescent idiopathic scoliosis: a randomised controlled clinical trial. *Clin Rehabil*. 2016 Feb;30(2):181-90.
41. Liu D, Yang Y, Yu X, et al. Effects of Specific Exercise Therapy on Adolescent Patients With Idiopathic Scoliosis: A Prospective Controlled Cohort Study. *Spine (Phila Pa 1976)*. 2020 Aug 1;45(15):1039-1046.
42. Schreiber S, Parent EC, Moez EK, et al. The effect of Schroth exercises added to the standard of care on the quality of life and muscle endurance in adolescents with idiopathic scoliosis-an assessor and statistician blinded randomized controlled trial: "SOSORT 2015 Award Winner," *Scoliosis*. 2015 Sep 18;10:24.
43. Schreiber S, Parent EC, Khodayari Moez E, et al. Schroth Physiotherapeutic Scoliosis-Specific Exercises Added to the Standard of Care Lead to Better Cobb Angle Outcomes in Adolescents with Idiopathic Scoliosis - an Assessor and Statistician Blinded Randomized Controlled Trial. *PLoS One*. 2016 Dec 29;11(12):e0168746.
44. Schreiber S, Parent EC, Hill DL, et al. Schroth physiotherapeutic scoliosis-specific exercises for adolescent idiopathic scoliosis: how many patients require treatment to prevent one deterioration? - results from a randomized controlled trial - "SOSORT 2017 Award Winner". *Scoliosis Spinal Disord*. 2017 Nov 14;12:26.
45. Schreiber S, Parent EC, Hill DL, et al. Patients with adolescent idiopathic scoliosis perceive positive improvements regardless of change in the Cobb angle - Results from a randomized controlled trial comparing a 6-month Schroth intervention added to standard care and standard care alone. *SOSORT 2018 Award winner*. *BMC Musculoskelet Disord*. 2019 Jul 8;20(1):319.
46. Gao C, Zheng Y, Fan C, et al. Could the clinical effectiveness be improved under the integration of orthotic intervention and scoliosis-specific exercise in managing adolescent idiopathic scoliosis? *Am J Phys Med Rehabil*. 2019;98:642-648.
47. Simhon ME, Fields MW, Grimes KE, et al. Completion of a formal physiotherapeutic scoliosis-specific exercise training program for adolescent idiopathic scoliosis increases patient compliance to home exercise programs. *Spine Deform*. 2021;9(3):691-696.
48. Fan Y, Ren Q, To MKT, Cheung JPY. Effectiveness of scoliosis-specific exercises for alleviating adolescent idiopathic scoliosis: A systematic review. *BMC Musculoskelet Disord*. 2020;21(1).