





























































- Upasani VV, Newton PO. Ten-Year Outcomes of Selective Fusions for Adolescent Idiopathic Scoliosis. *J Bone Joint Surg Am.* 2019 May 1;101(9):761-770.
16. Shah SA, Dhawale AA, Oda JE, Yorgova P, Neiss GI, Holmes L Jr, Gabos PG. Ponte Osteotomies With Pedicle Screw Instrumentation in the Treatment of Adolescent Idiopathic Scoliosis. *Spine Deform.* 2013 May;1(3):196-204.
17. Holewijn RM, Schlösser TP, Bisschop A, van der Veen AJ, Stadhouders A, van Royen BJ, Castelein RM, de Kleuver M. How Does Spinal Release and Ponte Osteotomy Improve Spinal Flexibility? The Law of Diminishing Returns. *Spine Deform.* 2015 Sep;3(5):489-495.
18. Halanski MA, Cassidy JA. Do multilevel Ponte osteotomies in thoracic idiopathic scoliosis surgery improve curve correction and restore thoracic kyphosis? *J Spinal Disord Tech.* 2013 Jul;26(5):252-5.
19. Koerner JD, Patel A, Zhao C, Schoenberg C, Mishra A, Vives MJ, Sabharwal S. Blood loss during posterior spinal fusion for adolescent idiopathic scoliosis. *Spine (Phila Pa 1976).* 2014 Aug 15;39(18):1479-87.
20. Samdani AF, Bennett JT, Singla AR, Marks MC, Pahys JM, Lonner BS, Miyanji F, Shah SA, Shufflebarger HL, Newton PO, Asghar J, Betz RR, Cahill PJ. Do Ponte Osteotomies Enhance Correction in Adolescent Idiopathic Scoliosis? An Analysis of 191 Lenke 1A and 1B Curves. *Spine Deform.* 2015 Sep;3(5):483-488.
21. Floccari, L. Ponte Osteotomies in a Matched Series of Large AIS Curves Increase Surgical Risk Without Improving Outcomes in Pediatric Orthopaedic Society of North America. 2019.
22. Goobie SM, Zurakowski D, Glotzbecker MP, McCann ME, Hedequist D, Brustowicz RM, Sethna NF, Karlin LI, Emans JB, Hresko MT. Tranexamic Acid Is Efficacious at Decreasing the Rate of Blood Loss in Adolescent Scoliosis Surgery: A Randomized Placebo-Controlled Trial. *J Bone Joint Surg Am.* 2018 Dec 5;100(23):2024-2032.
23. Bartley CE, Bastrom TP, Newton PO. Blood Loss Reduction During Surgical Correction of Adolescent Idiopathic Scoliosis Utilizing an Ultrasonic Bone Scalpel. *Spine Deform.* 2014 Jul;2(4):285-290.
24. Murgai RR, Andras LM, Nielsen E, Scott G, Gould H, Skaggs DL. Dedicated spine nurses and scrub technicians improve intraoperative efficiency of surgery for adolescent idiopathic scoliosis. *Spine Deform.* 2020 Apr;8(2):171-176.
25. Borden TC, Bellaire LL, Fletcher ND. Improving perioperative care for adolescent idiopathic scoliosis patients: the impact of a multidisciplinary care approach. *J Multidiscip Healthc.* 2016 Sep 14;9:435-445.
26. Mankin KP, Moore CA, Miller LE, Block JE. Hemostasis with a bipolar sealer during surgical correction of adolescent idiopathic scoliosis. *J Spinal Disord Tech.* 2012 Jul;25(5):259-63.
27. Andras L, Yamaguchi KT Jr, Skaggs DL, Tolo VT. Surgical technique for balancing posterior spinal fusions to the pelvis using the T square of Tolo. *J Pediatr Orthop.* 2012 Dec;32(8):e63-6.
28. Choudhry DK, Brenn BR, Sacks K, Shah S. Evaluation of Gabapentin and Clonidine Use in Children Following Spinal Fusion Surgery for Idiopathic Scoliosis: A Retrospective Review. *J Pediatr Orthop.* 2019 Oct;39(9):e687-e693.
29. Shau DN, Bible JE, Gadowski SP, Samade R, Armaghani S, Mencio GA, Devin CJ. Utility of Postoperative Radiographs for Pediatric Scoliosis: Association Between History and Physical Examination Findings and Radiographic Findings. *J Bone Joint Surg Am.* 2014 Jul 2;96(13):1127-1134.